

Content available at: <https://www.ipinnovative.com/open-access-journals>

Journal of Preventive Medicine and Holistic Health

Journal homepage: <https://www.jpmmh.org/>

Case Report

A case report of successful treatment approach for Vaginismus

Rozina Shaikh¹, Binal Dave^{1,*}

¹Dept. of Women's Healthcare, Aqua Centric Private Limited, Mumbai, Maharashtra, India



ARTICLE INFO

Article history:

Received 29-12-2021

Accepted 08-01-2022

Available online 30-06-2022

Keywords:

Dyspareunia

Pelvic Floor Dysfunction

Hypertonic pelvic floor dysfunction

Pelvic floor rehab

ABSTRACT

Vaginismus is a condition that can be defined as an uncontrolled contraction of the vaginal muscles, which can lead to difficulty in coital activity.^{1,2} The term vaginismus was coined in 19th century. However, vaginismus has been actualized as a inconsistent but well recognized and well managed female sexual dysfunction. In 1859 gynecologist pen down from his personal experience “I can confidently assert that I know of no disease capable of producing so much unhappiness to both parties of the marriage contract, and I am happy to state that I know of no serious trouble that can be cured so easily, so safely and so certainly”^{1,3}. This actualization was extended by Masters and Johnson. Who clocked in a treatment and outcome success rate of 100%.^{1,4} Beck stated vaginismus as “an interesting illustration of scientific neglect”^{1,5}.

There is paucity in the evidences of epidemiological studies examining the population prevalence of vaginismus as it requires gynecological assessment and the effected will avoid it due the pain anticipation due to which there have been numerous estimation with concerns to the prevalence vaginismus.¹ Masters and Johnson state that it is comparatively sparse condition^{1,4,6} And there are others who advocate that it is the most found female psychosexual dysfunction.⁷⁻¹⁰ The prevalence of vaginismus remains unspecific however the prevalence rate in clinical setting have been outlined to fall between 5-17%¹¹.

It was found that Ghana reported 68.1% cases with signs and symptoms of vaginismus.¹² An Italian women reported 9% of enquires for vaginismus over a phone call helpline.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Vaginismus is a condition that can be defined as an uncontrolled contraction of the vaginal muscles, which can lead to difficulty in coital activity.^{1,2} The term vaginismus was coined in 19th century. However, vaginismus has been actualized as a inconsistent but well recognized and well managed female sexual dysfunction. In 1859 gynecologist pen down from his personal experience “I can confidently assert that I know of no disease capable of producing so much unhappiness to both parties of the marriage contract, and I am happy to state that I know of no serious trouble that

can be cured so easily, so safely and so certainly”.¹⁻³ This actualization was extended by Masters and Johnson. Who clocked in a treatment and outcome success rate of 100%.^{1,4} Beck stated vaginismus as “an interesting illustration of scientific neglect”^{1,5}

There is paucity in the evidences of epidemiological studies examining the population prevalence of vaginismus as it requires gynecological assessment and the effected will avoid it due the pain anticipation due to which there have been numerous estimation with concerns to the prevalence vaginismus.¹ Masters and Johnson state that it is comparatively sparse condition^{1,4,6} And there are others who advocate that it is the most found female psychosexual dysfunction.⁷⁻¹⁰ The prevalence of vaginismus remains

* Corresponding author.

E-mail address: Doctorbinal85@gmail.com (B. Dave).

unspecific however the prevalence rate in clinical setting have been outlined to fall between 5-17%.¹¹

It was found that Ghana reported 68.1% cases with signs and symptoms of vaginismus.¹² An Italian women reported 9% of enquires for vaginismus over a phone call helpline for sexual problems.¹³ Women in Iran who attended a family planning clinic reported 12% of women suffered from vaginismus at least 50% of the time with 4% suffering vaginismus.¹⁴ A study in Turkey where 54 women attended the psychiatric OPD found 75.9% reported vaginismus out of which 6 turned up with a lifelong problem¹⁵

2. Need of The Study

Vaginismus (Dyspareunia) being very commonly seen in women but rarely spoken about making it difficult to target the population. The need of the study is to create awareness about vaginismus and possible physiotherapy management for the same.

3. Case Report

1. A 29 years old women nulliparous IT professional with no medical history was referred to us with a chief complaint of pain during coital activity and unable to complete the coital activity due to pain. When the comprehensive analysis was done her BMI was 20.3kg/m² she had an increased lumbar lordosis and her muscle strength according to manual muscle testing was as follows
2. Lower abdominal strength (Transverse abdominis : 3/5)
3. Int and Ext Obliques: 3/5
4. Lower back strength (Multifidus: 3+/5)
5. DRAM: supraumbilical - 2 fingers
6. Umbilical - 2 fingers
7. Infraumbilical - 2 $\frac{1}{2}$ fingers

Transvaginal examination findings are in Box 1 with states the finding done on the day of consultation and on her 1st review

4. Pelvic Floor Physiotherapy Intervention

4.1. Relaxation

Breathing relaxation was given to the women where the 1st 10 mins of the intervention was invested in. she was asked to take a deep breath in from the nose and breathe out from the mouth once this concept was familiarized then she was asked to breath in from the nose and breath out from the mouth and at the same time try to open the virginal opening.

4.2. Perineal massage

For the the individual was asked to lie down on her back with knees folded then she was asked to rest her knees on the therapist. Gentle strokes were given at the vaginal opening

Table 1:

Measurments	Day 1-01/12/20	1 st review -04/02/2021
Dynamic assessment	Accessory muscles use	No accessory muscles use
Response to cough	Inward Protrusion	Inward Protrusion
Palpation	Outer ring tighter than the inner ring (1finger penetration	Outer ring comparatively less tight (allowed 1 finger penetration), trigger point at 3,9,10 o'clock
PFm Relaxtion (HET's MMT)	-1	-2 on the 1 st 2 tries -3 from 3 rd tries
Endurance	5 Sec	8 Sec
Repetition	4	7
Fast twitch	4	6

Table 2:

Out come measures	Day 01-01/12/20	1 st review-04/02/2021
Dilator size	1 finger penetration was difficulty	Dilator size 4 dilator was introduced with a little difficulty but penetration was possible
Female sexual function index (FSFI)	6.6	24.6

and strokes from mons pubis to vaginal opening with gloved fingers and gel applied on the fingers. Gentle pressure with the tips of the fingers was applied at the vaginal opening.

4.3. Clock wise awareness of pelvic floor

Once she was comfortable with the relaxation and the perineal massage clockwise awareness was thought to her this would make it easy for her to understand her pelvic floor and understand where the tightness is.

4.4. Finger insertion

After the pelvic awareness, gently a finger is inserted into the vaginal opening this is done by keeping the gloved finger at the vaginal opening and asking her to contract and relax. This continuous contraction and relaxation will guide the finger into the vaginal canal without applying excessive forces and stimulating pain. Once this is done with less pain and minimal discomfort, it will help in gaining her confidence in you and will also help in reducing her anxiety regarding penetration. Once the finger in the vaginal canal very gently the clockwise assessment was done. The finding of the assessment on the 1st day and the review is in box 1.

4.5. Trigger point release

After the clockwise assessment and the trigger points are determined they are gently released by applying pressure as tolerated by her and the pressure is maintained for 30 sec. this is repeated 3 to 4 times and then the other trigger point is targeted.

4.6. Dilator

After the trigger points are released and the finger penetration gets easier with lesser pain dilator was introduced. Use of dilator must be in functional positioning. Use of the dilator was started in lying down position, once this was achieved was progressed to side lying and then into all 4 position. The dilator was first applied to the applicator and with gel applied at the tip and girth of the dilator was gently placed at the vaginal canal and with the same technique that was used for finger penetration the dilator was introduced into the vaginal canal. Once the dilator was introduced into the vaginal cavity gently the copulation activity was mimicked and was continued for 15-20 mins with rest period of 2 3 mins after every 4 mins.

5. Discussion

The invasive therapy used for managing the case shows a positive outcome to the therapy delivered. There is an improvement seen in the hypertonus muscle and a significant improvement in the quality of life. In a study done by Weiss JM it states that invasive pelvic myofascial release helps in reducing the hypertonus state of the pelvic floor muscle¹⁶ similarly in the case study performed the invasive intervention given for the release of the trigger points in the pelvic has shown to reduce the hypertonicity and was maintained till the next session.

There was a study performed by Silva AP Et al. where perineal massage was done for individuals with dyspareunia and showed positive effects in reducing dyspareunia caused by tenderness¹⁷ similarly in this case the tenderness was highly prevalent and very painful and was building the anxiety and fear in the individual which was similar to the study done by Payne KA et al.¹⁸ perineal massage helped in reducing the tenderness, pain and the anxiety which was associated with pain which helped in improving penetration.

In a study by Pacik PT et al dilators were used in a case of vaginismus which helped in improving and maintaining the stretch¹⁹ similarly in this case the dilator helped in stretching and maintaining the tone of the pelvic floor muscle which helped in improving the quality of penetration.

In a study conducted by C M Meston on the scoring of the FSFI has the weight to correctly understand the sexual dysfunction and to a guided treatment.²⁰

6. Conclusion

The treatment helped in reducing the tone of the pelvic floor, reduced the intensity of trigger points, reduced the anxiety and fear of penetration resulting in improved quality of copulation.

7. Source of Funding

None.

8. Conflict of Interest

None.

References


- Lahaie MA, Boyer SC, Amsel R, Khalifé S, Binik YM. Vaginismus: a review of the literature on the classification/diagnosis, etiology and treatment. *Women's Health*. 2010;6(5):705–24. doi:10.2217/whe.10.46.
- American Psychiatric Association: Diagnostic and Statistical Manual of Mental Disorders.; 2000. p. 992. Available from: http://repository.poltekkes-kaltim.ac.id/6571/Diagnostic%20and%20statistical%20manual%20of%20mental%20disorders%20_%20DSM-5%20%28%20PDFDrive.com%20%29.pdf.
- Sims MJ. On vaginismus. *Trans Obstet Soc London*. 1861;3(1):356–67.
- Masters WH, Johnson VE. Human sexual inadequacy. By W. H. Masters and V. E. Johnson. Little, Brown, Boston. . *Teratology*. 1970;4(4):465–7. doi:10.1002/tera.1420040411.
- Beck J, Vaginismus. Handbook of Sexual Dysfunctions: Assessment and Treatment. W O, JH G, editors. Boston, USA: Allyn and Bacon Inc; 1993. p. 381–397.
- Schmidt G, Arentewicz G. The Treatment of Sexual Disorders: Concepts and Techniques of Couple Therapy; 1982. p. 123–46.
- Simons JS, Carey MP. Prevalence of sexual dysfunctions: results from a decade of research. *Arch Sex Behav*. 2001;30(2):177–219. doi:10.1023/a:1002729318254.
- Crowley T, Richardson D, Goldmeier D. Recommendations for the management of vaginismus: BASHH special interest group for sexual dysfunction. *Int J STD AIDS*. 2006;17(1):14–8. doi:10.1258/095646206775220586.
- Kabakçi E, Batur S. Who bene ts from cognitive behavioural therapy for vaginismus? *J Sex Marital Ther*. 2003;29(4):277–88. doi:10.1080/00926230390195515.
- Mcguire H, Hawton K. Interventions for vaginismus. *Cochrane Database Syst Rev*. 2001;12:1760. doi:10.1002/14651858.CD001760.pub2.
- Spector I, Carey M. Incidence and prevalence of the sexual dysfunctions: a critical review of the empirical literature. *Arch Sex Behav*. 1990;19(4):389–96. doi:10.1007/BF01541933.
- Amidu N, Owiredu W, Woo E, Mensah OA, Quaye L, Alhassan A. Incidence of sexual dysfunction: a prospective survey in Ghanaian females. *Reprod Biol Endocrinol*. 2010;8(1):106–17. doi:10.24966/RMGO-2574/100046.
- Simonelli C, Tripodi F, Cosmi V, Rossi R, Fabrizi A, Silvaggi C. What do men and women ask a helpline on sexual concerns? Results of an Italian telephone counselling service. *Int J Clin Pract*. 2010;64(3):360–70. doi:10.1111/j.1742-1241.2009.02269.x.
- Shokrollahi P, Mirmohamadi M, Mehrabi F, Babaei GH. Prevalence of sexual dysfunction in women seeking services at a family planning centers in Tehran. *J Sex Marital Ther*. 1999;25(3):211–6. doi:10.1080/00926239908403995.
- Dogan S. Vaginismus and accompanying sexual dysfunctions in a Turkish clinical sample. *J Sex Med*. 2009;6(1):184–92. doi:10.1111/j.1743-6109.2008.01048.x.

16. Weiss JM. Pelvic floor myofascial trigger points: manual therapy for interstitial cystitis and the urgency-frequency syndrome. *J Urol.* 2001;166(6):2226–57. doi:10.1016/s0022-5347(05)65539-5.
17. Silva AP, Montenegro ML, Gurian MB, Mitidieri AM, Lara LA, Poli OB. Perineal massage improves the dyspareunia caused by tenderness of the pelvic floor muscles. *Rev Bras Ginecol Obstet.* 2017;39(1):26–30. doi:10.1055/s-0036-1597651.
18. Payne KA, Binik YM, Amsel R, Khalifé S. When sex hurts, anxiety and fear orient attention towards pain. *Eur J Pain.* 2005;9(4):427–63. doi:10.1016/j.ejpain.2004.10.003.
19. Pacik PT, Geletta S. Vaginismus treatment: clinical trials follow up 241 patients. *Sexual Med.* 2017;5(2):114–37. doi:10.1016/j.esxm.2017.02.002.
20. Meston CM, Freihart BK, Handy AB, Kilimnik CD, Rosen RC. Scoring and Interpretation of the FSFI: What can be Learned From 20 Years of use?. *J Sex Med.* 2020;17(1):17–25.

doi:10.1016/j.jsxm.2019.10.007.

Author biography

Rozina Shaikh, Womens Health Care Therapist

Binal Dave, Lead Orthopaedic and Womens Health care
 <https://orcid.org/0000-0001-8512-5627>

Cite this article: Shaikh R, Dave B. A case report of successful treatment approach for Vaginismus. *J Prev Med Holistic Health* 2022;8(1):42-45.