



Content available at: <https://www.ipinnovative.com/open-access-journals>

Journal of Preventive Medicine and Holistic Health

Journal homepage: <https://www.jpmmh.org/>



Original Research Article

Efficiency of rehabilitation in limb asymmetries of achilles tendinopathy female patients

Mohammed Sheeba Kauser^{1,*}, Subhasis Karmakar², Laxmi Devi³, Irfan Ullah⁴

¹AR Clinic, Andhra Pradesh, India

²Private Practitioner, United Arab Emirates

³Parul University, Vadodara, Gujarat, India

⁴NF Railway Sports Association, Guwahati, Assam, India



ARTICLE INFO

Article history:

Received 20-05-2023

Accepted 06-06-2023

Available online 07-06-2023

Keywords:

Achilles tendinopathy

Between appendage unevenness

Weighty sluggish opposition

preparing

Highload

Physiotherapy

Ligament

ABSTRACT

Background: We want to learn more about heel pain in patients since they have been linked to increased injury risk.

Materials and Methods: We evaluated the effects of a 8-week exercise intervention mechanical, material, and morphological musculoskeletal characteristics and function in patients with heel pain (n = 44). Additionally, we investigated the relationship between asymmetry reductions and better patient-reported outcomes.

Results: At baseline, tendons with tendinopathy showed decreased tendon force (p = 0.013), decreased tendon stress (p 0.0001), greater tendon cross-sectional area (p 0.001),.

Conclusion: The injured leg contrasts from the asymptomatic leg while helpful activity mediations meaningfully affect imbalances.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Ankle pain is one of the most widely recognized wounds of the lower limbs. It is described by changes of ligament microstructure and is related with torment, enlarging, furthermore, a deficiency of capability¹ Albeit multifactorial in beginning, ligament mechanical overburden is viewed as the super contributing element.²⁻⁶

Between deviation might add to either the beginning or the ingenuity of musculoskeletal injury. Past exploration revealed that a between ankle pain is more noteworthy than 10-15% is related with expanded injury risk and diminished execution,^{7,8} while between imbalances lower than 10% for strength or capability are considered as threshold permitting get back to-wear.⁹ With regards to calcaneal injuries more prominent has been identified in patellar tendinopathy patients in contrast with controls was joined by a higher lower appendage outer muscle injury

* Corresponding author.

E-mail address: sheebaishaq.doc@gmail.com (M. Sheeba Kauser).

risk in everyday in those patients.¹⁰ Between appendage strength deviation might prompt non-symmetric stacking, which may thus expand the gamble of over-burdening,¹¹ inconsistent power assimilation, and decreased security¹² because of changes in engine conduct.¹³ To be sure, strength deviation has been accounted for to be a gamble factor for lower limb injury pain.,¹⁴ recommending a negative effect of articulated between appendage contrasts.

2. Methodology and Results

The Alfredson team females performed erratic activities with six arrangements of 15 redundancies two times everyday as indicated by the usually known convention.^{15–20} According to convention, flighty as it were withdrawals of the plantar flexors of the impacted leg were performed (heel-drop) while the sound leg performed concentric plantar flexor muscle withdrawals (heel-ascend) to get back to the beginning position. The high-load bunch played out a profoundly serious one-sided strength

preparing convention comprising of five arrangements of four plantar flexor withdrawals at 90% of isometric greatest wilful constrictions held for three seconds followed by three seconds of rest each and with 1-min in the middle between sets with the impacted leg.^{21–30}

This convention was executed four times each week with a portable input fitted sling which has recently been depicted.³¹ The dissemination of meetings in the span of seven days was picked independently. Load movement was took into account the two gatherings after the initial fourteen days, if torment level on the numeric rating (NRS) scale was $<4/10$ ³² and the singular rating seen effort was $<5/10$ (NRS scale). Because of moral reasons, all patients gotten the choice of 12 meetings of latent restorative treatment (i.e., manual treatment, tissue, or joint preparation). Besides, members were permitted to go on with their preparing propensities in the event that aggravation levels were $\leq 5/10$ (NRS scale). Any extra strength preparing of the plantar flexors or recently began lower body preparing was restricted during the intervention stage. This was compared with the normal group B which were with traditional therapy, ultrasound and simple RICE therapy.(Table 1)

3. Discussion

In concurrence with the writing and with our speculation, we found between imbalances in Ankle injury .While looking at the asymptomatic side, the ligaments was portrayed by huge lower ligament force, lower ligament stress,.

While a specific level of limb imbalances can be viewed as non-obsessive, articulated between leg contrasts especially in ligament and vascularization have all the earmarks of being a sign of ankle injury. Augmented tendinopathic ligaments have been accounted for in many examinations^{33–35} and might be brought about by irritation related edema. Restricted fusiform ligament enlarging has been accounted for already in midportion tendinopathy. Nonetheless, our discoveries demonstrate that the expansion in ligament was not confined to a particular area inside the ligament yet was evident all through the whole length of the free. In spite of the fact that leg predominance would be able influence between appendage ligament, laterality was not connected with the side of injury in our review, making it improbable was brought about by more serious stacking.^{36–40}

Conversely, ligament mechanical properties like decreased solidness or expanded strain have been examined to be a gamble factor for tendinopathy. Flow research evaluating between leg imbalances for mechanical ligament properties in tendinopathy revealed decreased ligament strain and diminished ligament firmness. Moreover, in light of correlations with matched controls, lower ligament firmness and higher ligament strain have been accounted for in calcaneal tendinopathy. Shear wave elastography estimations affirmed discoveries of diminished ligament firmness, prompting the possibility of solidness measures as an expected marker for tendinopathic changes in a new survey. By the by, we didn't identify any huge between appendage ligament firmness nor ligament strain contrasts.

Thus, a more equilibrated ligament strain between appendages could safeguard the tendinopathic ligament against strain-incited re-injury. In addition, our examinations of the PRE to POST contrasts between the two methods showed a reduction for the AAI of ligament.^{40–43}

Table 1:

	Group A.	Group B
Age.	35 - 45 years	33 - 46 years
Sex	Females	Females
Height.	131 - 165 cm.	136 - 164 cm
Weight.	45 to 70 kgs.	47 to 75 kgs
Pre test	3.5 -/+ 2.2	4.5+/- 6.5
Post test (p value).	0.03	0.018

Thus, a decrease in unevenness can't really be deciphered as being better. Albeit the two sorts of activity mediation decreased tendinopathic side effects an improvement in ligament wellbeing was essentially connected with in two factors as it were. Those two factors varied between gatherings and are subsequently not delegate for a general impact. Besides, We have previously detailed that three months of one-sided high-stacking practice in tendinopathy patients expanded ligament firmness the prepared leg.

In solid ligaments, it is deeply grounded that high-stacking exercise conventions prompting bigger ligament strains bring about higher expansions in mechanical and morphological ligament properties than low-strain conventions. This was as of late affirmed in tendinopathy patients.³⁵

In such manner, POST-to-POST contrasts between the group a and b team showing the high-load bunch in ligament strain what's more, ligament prolongation are in accordance with the writing essentially to the degree that an effect on ligament strain and along these lines ligament prolongation may be normal. Thus, a more equilibrated ligament strain between appendages could safeguard the tendinopathic ligament against strain-incited re-injury. In addition, our examinations of the PRE to POST contrasts between the two gatherings showed a reduction for the AAI of ligament. In such manner, outstanding unpredictable activity convention prompts unique stacking of the two appendages, as each redundancy comprises of a concentric plantar flexion (heel rise) of the unaffected appendage and a whimsical dorsal flexion (heel drop) of the impacted appendage. one-sided practices and may can possibly decrease between appendage imbalance in reciprocal developments like the CMJ.

While the two-sided nature of the activity might make sense of why VISA-A enhancements related with an AAI

decrease of CMJ percent drive proportion in this gathering, the erratic and concentric activities might be related with contrasts in ligament stacking which might have caused the AAI increment of ligament firmness recognized in this gathering.

4. Limitation

The unaffected appendage ought to be viewed as "ordinary" since proof recommends control changes to happen in the two legs. As this study has been led with female patients just the exchange of the outcomes to male patients is restricted, especially as we probably are aware that side effects and regenerative cycles might vary between genders.

Future exploration could additionally research the almost negligible difference among physiological and neurotic to lay out a more significant comprehension of relative between appendage unevenness changes and their clinical ramifications.

5. Conclusion

To illuminate clinical practice, choices with respect to-game should neither solely be founded on the presence nor the decrease of between appendage imbalance. In such manner, it very well may be profitable to procure standard pre-injury information for the two appendages in competitors which can be utilized as reference in the event of injury. Future exploration could additionally research the almost negligible difference among physiological and neurotic to lay out. In tendinopathy the suggestive side altogether contrasts from the asymptomatic contralateral appendage. Notwithstanding, the job and importance of unevenness with respect to ligament wellbeing is as yet muddled as, with the exception of ligament vascularity, the AAIs are in a similar reach as recently found in the solid populace. While

explicit one-sided or reciprocal activity intercessions may have little yet contrasting impacts on deviations, a decrease in imbalance isn't necessarily related with an improvement in ligament wellbeing while an expansion in unevenness doesn't be guaranteed to show a decrease in wellbeing.

6. Source of Funding

None.

7. Conflict of Interest


None.

References

- Riley G. Tendinopathy-from basic science to treatment. *Nat Clin Pract Rheumatol*. 2008;4(2):82–9.
- Lin TW, Cardenas L, Soslowky LJ. Biomechanics of tendon injury and repair. *J Biomech*. 2004;37(6):865–77.
- Magnusson SP, Langberg H, Kjaer M. The pathogenesis of tendinopathy: balancing the response to loading. *Nat Rev Rheumatol*. 2010;6(5):262–8.
- September A, Rahim M, Collins M. Towards an understanding of the genetics of tendinopathy. *Adv Exp Med Biol*. 2016;920:109–16. doi:10.1007/978-3-319-33943-6_9.
- Hopkins C. This paper highlights the socioeconomic importance of tendinopathy in medical practice. *Asia Pac J Sports Med Arthrosc Rehabil Technol*. 2016;4:9–20.
- Albers IS, Zwerver J, Diercks RL, Dekker JH, Van den Akker-Scheek, I. Incidence and prevalence of lower extremity tendinopathy in a Dutch general practice population: a cross sectional study. *BMC Musculoskelet Disord*. 2016;17:16. doi:10.1186/s12891-016-0885-2.
- Riel H, Lindstrom CF, Rathleff MS, Jensen MB, Olesen JL. Prevalence and incidence rate of lower-extremity tendinopathies in a Danish general practice: a registry-based study. *BMC Musculoskelet Disord*. 2019;20(1):239. doi:10.1186/s12891-019-2629-6.
- Linsell L. Prevalence and incidence of adults consulting for shoulder conditions in UK primary care: patterns of diagnosis and referral. *Rheumatology*. 2006;45(2):215–21.
- Windt DD, Koes BW, De Jong B, Bouter LM. Shoulder disorders in general practice: incidence, patient characteristics, and management. *Ann Rheum Dis*. 1995;54(12):959–64.
- O'neill S, Barry S, Watson P. Plantarflexor strength and endurance deficits associated with mid-portion Achilles tendinopathy: The role of soleus. *Phys Ther Sport*. 2019;37:69–76.
- Kujala UM, Sarna S, Kaprio J. Cumulative incidence of achilles tendon rupture and tendinopathy in male former elite athletes. *Clin J Sports Med*. 2005;15(3):133–5.
- Kader D, Saxena A, Movin T, Maffulli N. Achilles tendinopathy: Some aspects of basic science and clinical management. *Br J Sports Med*. 2002;36(4):239–49.
- Abate M, Silbernagel KG, Siljeholm C, Iorio AD, De Amicis D, Salini V. Pathogenesis of tendinopathies: Inflammation or degeneration? *Arthritis Res Ther*. 2009;11(3):235. doi:10.1186/ar2723.
- Sharma P, Maffulli N. Biology of tendon injury: Healing, modeling and remodeling. *J Musculoskelet Neuronal Interact*. 2006;6(2):181–90.
- Schmitt LC, Paterno MV, Hewett TE. The Impact of Quadriceps Femoris Strength Asymmetry on Functional Performance at Return to Sport Following Anterior Cruciate Ligament Reconstruction. *J Orthop Sports Phys Ther*. 2012;42(9):750–9.
- Fendri T, Rebai H, Harrabi MA, Chaari F, Boyas S, Beaune B. Athletes with unilateral patellar tendinopathy have increased subsequent lower extremity musculoskeletal injury risk. *Eur J Sports Sci*. 1908;22(12):1908–15.
- Ford KR, Myer GD, Hewett TE. Valgus knee motion during landing in high school female and male basketball players. *Med Sci Sports Exerc*. 2003;35(10):1745–50.
- Paterno MV, Schmitt LC, Ford KR, Rauh MJ, Myer GD, Huang B. Biomechanical measures during landing and postural stability predict second anterior cruciate ligament injury after anterior cruciate ligament reconstruction and return to sport. *Am J Sports Med*. 1968;38(10):1968–78.
- Helme M, Tee J, Emmonds S, Low C. Does lower-limb asymmetry increase injury risk in sport? A systematic review. *Phys Ther Sport*. 2021;49:204–13. doi:10.1016/j.ptsp.2021.03.001.
- Bains BS, Porter K. Lower limb tendinopathy in athletes. *Trauma*. 2006;8(4):213–24.
- Mcauliffe S, Tabuena A, McCreesh K, O'keeffe M, Hurley J, Comyns T, et al. Altered Strength Profile in Achilles Tendinopathy: A Systematic Review and Meta-Analysis. *J Athl Train*. 2019;54(8):889–900.
- Hasani F, Vallance P, Haines T, Munteanu SE, Malliaras P. Are Plantarflexor Muscle Impairments Present Among Individuals with Achilles Tendinopathy and Do They Change with Exercise? A Systematic Review with Meta-analysis. *Sports Med Open*. 2021;7(1):18. doi:10.1186/s40798-021-00308-8.
- Chimenti RL, Flemister AS, Tome J, McMahan JM, Flannery MA, Xue Y. Altered tendon characteristics and mechanical properties associated with insertional achilles tendinopathy. *J Orthop Sports Phys Ther*. 2014;44(9):680–9.
- Tillander B, Gauffin H, Lyth J, Knutsson A, Timpka T. Symptomatic Achilles Tendons are Thicker than Asymptomatic Tendons on Ultrasound Examination in Recreational Long-Distance Runners. *Sports*. 2019;7(12):245. doi:10.3390/sports7120245.
- Hirschmüller A, Frey V, Deibert P, Konstantinidis L, Mayer F, Sudkamp N. Achilles tendon power Doppler sonography in 953 long distance runners-A cross sectional study. *Ultraschall Med*. 2010;31(4):387–93.
- Crawford SK, Thelen D, Yakey JM, Heiderscheid BC, Wilson JJ, Lee KS. Regional shear wave elastography of Achilles tendinopathy in symptomatic versus contralateral Achilles tendons. *Eur Radiol*. 2023;33(1):720–9.

27. Risch L, Stoll J, Schomöller A, Engel T, Mayer F, Cassel M. Intra-individual Doppler Flow Response to Exercise Differs Between Symptomatic and Asymptomatic Achilles Tendons. *Front Physiol.* 2021;12:617497. doi:10.3389/fphys.2021.617497.
28. Agres AN, Duda GN, Gehlen TJ, Arampatzis A, Taylor WR, Manegold S, et al. Increased unilateral tendon stiffness and its effect on gait 2–6 years after Achilles tendon rupture. *Scand. J Med Sci Sports.* 2015;25(6):860–7.
29. Sánchez AB, Abián P, Jimenez F. Structural and mechanical properties of the Achilles tendon in senior badminton players: Operated vs. non-injured tendons. *Clin Biomech.* 2021;85:105366. doi:10.1016/j.clinbiomech.2021.105366.
30. Hoeffner R, Svensson RB, Bjerregaard N, Kjær M, Magnusson SP. Persistent Deficits after an Achilles Tendon Rupture: A Narrative Review. *Transl Sports Med.* 2022;7445398. doi:10.1155/2022/7445398.
31. Bohm S, Mersmann F, Marzilger R, Schroll A, Arampatzis A. Asymmetry of Achilles tendon mechanical and morphological properties between both legs. *J Med Sci Sports.* 2015;25(1):124–32.
32. Sánchez AB, Abián P, Jiménez F, Vicén JA, Bayliss AJ, Weatherholt AM, et al. Myotendinous asymmetries derived from the prolonged practice of badminton in professional players. *J Musculoskeletal Neuronal Interact.* 2016;14(9):105–12.
33. Bayliss AJ, Weatherholt AM, Crandall TT, Farmer DL, McConnell JC, Crossley KM. Achilles tendon material properties are greater in the jump leg of jumping athletes. *J Musculoskeletal Neuronal Interact.* 2016;16(2):105–12.
34. Hewett TE, Ford KR, Hoogenboom BJ, Myer GD. Understanding and preventing acl injuries: Current biomechanical and epidemiologic considerations-Update. *N Am J Sports Phys Ther.* 2010;5(4):234–51.
35. Paterno MV, Ford KR, Myer GD, Heyl R, Hewett TE. Limb asymmetries in landing and jumping 2 years following anterior cruciate ligament reconstruction. *Clin J Sports Med.* 2007;17(4):258–62. doi:10.1097/JSM.0b013e31804c77ea.
36. Rozzi SL, Lephart SM, Sterner R, Kuligowski L. Balance training for persons with functionally unstable ankles. *J Orthop Sports Phys Therapy.* 1999;29(8):478–86. doi:10.2519/jospt.1999.29.8.478.
37. Richie DR. Effects of foot orthoses on patients with chronic ankle instability. *J Am Podiatric.* 2007;97(1):19–30. doi:10.7547/0970019.
38. Shiravi Z, Talebian S, Hadian MR, Oliaie GR. Effect of cognitive task on postural control of the patients with chronic ankle sprain. *J Foot Ankle Res.* 2012;5(1):24.
39. Mckeon PO, Wikstrom EA. Sensory-targeted ankle rehabilitation strategies for chronic ankle instability. *Med Sci Sports Exerc.* 2016;48(5):776–84. doi:10.1249/MSS.0000000000000859.
40. Diaz DC, Vega RL, Pérez MSO, Contreras FH, Amat AM. Effects of 6 weeks of balance training on chronic ankle instability in athletes: a randomized controlled trial. *Int J Sports Med.* 2015;36(9):754–60. doi:10.1055/s-0034-1398645.
41. Cain MS, Ban RJ, Chen YP, Geil MD, Goerger BM, Linens SW, et al. Four-week ankle-rehabilitation programs in adolescent athletes with chronic ankle instability. *J Athletic Training.* 2020;55(8):801–10.
42. Alawna M, Mohamed AA. Short-term and long-term effects of ankle joint taping and bandaging on balance, proprioception and vertical jump among volleyball players with chronic ankle instability. *Physical Therapy in Sport.* 2020;46:145–54. doi:10.1016/j.ptsp.2020.08.015.
43. Tavakoli S, Forghany S, Nester C. The effect of dual tasking on foot kinematics in people with functional ankle instability. *Gait Posture.* 2016;49:364–70.

Author biography

Mohammed Sheeba Kauser, Chief Physiotherapist
 <https://orcid.org/0000-0001-5301-3559>

Subhasis Karmakar, Physiotherapist

Laxmi Devi, Phd Scholar

Irfan Ullah, Physiotherapist

Cite this article: Sheeba Kauser M, Karmakar S, Devi L, Ullah I. Efficiency of rehabilitation in limb asymmetries of achilles tendinopathy female patients. *J Prev Med Holistic Health* 2023;9(1):19-23.